

Quantum Field Theory in Curved Spacetime

Loso Judijanto¹, Bassam Al-Khouri², Rania Khatib³¹ IPOSS Jakarta, Indonesia² Mutah University, Jordan³ Jordan University of Science and Technology, Jordan

Corresponding Author:

Loso Judijanto,

IPOSS Jakarta, Indonesia

Jl. Kota Gedang No.28, RW.1, Ps. Manggis, Kecamatan Setiabudi, Kota Jakarta Selatan, Daerah Khusus Ibukota Jakarta 12970

Email: losojudijantobumn@gmail.com

Article Info

Received: March 10, 2025

Revised: June 8, 2025

Accepted: June 8, 2025

Online Version: June 8, 2025

Abstract

Quantum field theory and general relativity are the two main pillars of modern physics. However, the two still cannot be combined consistently to explain cosmic phenomena at the microscopic level, especially in the context of curved spacetime. This research aims to explore the interaction between quantum fields and the curvature of spacetime, with a focus on the implications of quantum gravity. This research aims to understand how quantum fields interact with curved spacetime, as well as to develop a more comprehensive model of physics that combines these two concepts. The methods used include the development of mathematical models and numerical simulations to integrate quantum field theory with general relativity. The analysis was carried out by examining the impact of space-time curvature on quantum field fluctuations around massive objects such as black holes. The findings show that the curvature of spacetime has a major influence on the behavior of the quantum field, leading to modifications in energy distribution and field fluctuations. This discovery opens up new possibilities in the development of a more complete theory of quantum gravity. This study provides new insights into understanding the relationship between quantum fields and curved spacetime, as well as opening the way for further research in the field of quantum gravity and extreme cosmic phenomena.

Keywords: Curved Spacetime, Quantum Field Theory, Quantum Gravity



© 2025 by the author(s)

This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution-ShareAlike 4.0 International (CC BY SA) license (<https://creativecommons.org/licenses/by-sa/4.0/>).

Journal Homepage

<https://journal.ypidathu.or.id/index.php/quantica>

How to cite:

Judijanto, L., Al-Khouri, B & Khatib, R. (2025). Quantum Field Theory in Curved Spacetime. *Journal of Tecnologia Quantica*, 2(1), 35–43.
<https://doi.org/10.70177/quantica.v2i1.1964>

Published by:

Yayasan Pendidikan Islam Daarut Thufulah

INTRODUCTION

Quantum field theory (QFT) is one of the main pillars in modern physics, which combines the principles of quantum mechanics with the theory of special relativity to describe the interactions between subatomic particles (Z. Wang et al., 2023). QFT provides an in-depth picture of how fields and particles interact on a microscopic scale (Tawfik & Dabash, 2023). The models in the QFT have successfully explained fundamental phenomena such as electromagnetic interactions, quantum gravity, and strong and weak nuclear forces.

In a flat spacetime, quantum field theory has managed to provide a very accurate description of physical phenomena (Mogull et al., 2021). However, the problem becomes more complicated when QFT is applied in the context of curved spacetime (Jakobsen et al., 2021). Albert Einstein developed a general theory of relativity that describes gravity as a space-time curvature caused by mass and energy (Hwang & Noh, 2023). Understanding how QFTs function in curved spacetime is a major challenge for theoretical physicists.

The theory of quantum fields in a flat spacetime has been confirmed through a number of experiments and observations (Liu et al., 2023). However, its application to curved spacetime requires a different approach due to the properties of gravity that alter the geometry of space and time (Kontou & Sanders, 2020). In this context, the quantum field must be considered as a phenomenon that is not only affected by the properties of the field itself, but also by the distortion of spacetime.

Cosmological physics and astrophysics, for example, show how quantum field theory interacts with gravity at very large and very small levels (G. Wang et al., 2023). The application of QFT in curved spacetime is important for understanding phenomena such as black holes, singularities, and other extreme events involving very strong gravitational influences (Bidussi et al., 2022). In this context, quantum gravity theory seeks to combine two different concepts, namely quantum theory and general relativity.

Recent developments in quantum gravity theory, such as string theory and loop quantum gravity theory, seek to provide a consistent framework for QFTs in curved spacetime (Wu & Zeng, 2022). These theories offer a new view of how particles and fields interact in dynamic spacetime geometry, although there is no consensus yet on which theory can fully explain the phenomenon.

The application of QFT in curved spacetime paves the way for further research in the fields of quantum gravity and cosmology (Adhikari et al., 2023). This involves more complex mathematical approaches and more in-depth experiments to confirm the predictions generated by these theories (Capolupo et al., 2020). A deeper understanding of the relationship between gravity and quantum fields could lead to new insights into the basic structure of the universe, and may lead to the discovery of a more fundamental theory that unifies all the laws of physics.

The theory of quantum fields in curved spacetime presents a major challenge that has not yet been fully solved in theoretical physics (Ra et al., 2020). Although many advances have been made in the incorporation of quantum mechanics and general relativity, there is not yet a consistent and fully understood framework for how these two theories interact at the microscopic scale when spacetime experiences curvature.

One area that has not been widely accepted is how quantum fields behave in curved spacetime. While QFTs in flat space can be precisely understood and predicted, their application to curved spacetime, especially around objects such as black holes or in early

universe conditions, presents great difficulties (Toscano-Negrette et al., 2023). This is due to the strong influence of gravity, which changes the geometry of space and time.

In the context of curved spacetime, the main problem lies in the merger between quantum and gravitational concepts (Kaubruegger et al., 2023). Gravity is not just a force, but a distortion in the structure of spacetime itself. Understanding how quantum fields interact with dynamic space-time geometry is a major gap in theoretical physics research that needs to be filled.

One of the main gaps that must be addressed is the mismatch between quantum field theory and general relativity in the face of extreme phenomena, such as gravitational singularities inside black holes (Maniccia et al., 2023). At this point, gravity and quantum effects must be explained in a consistent framework, but they cannot be combined without sacrificing mathematical and physical accuracy.

A limited understanding of the relationship between gravity and quantum in curved spacetime prevents us from reaching a theory of physics that holds together all fundamental interactions (Viermann et al., 2022). Therefore, understanding this gap is essential to create a theory that can explain the phenomena that occur at the smallest and largest levels in the universe.

Filling this gap is crucial for understanding the dynamics of spacetime in extreme conditions, such as near black holes or in the early universe (Shi et al., 2023). Quantum field theory applied to curved spacetime will allow us to better understand how quantum effects affect gravitational structures at the microscopic level. This opens up opportunities to explain phenomena that cannot be explained by current physical theories.

The merger between quantum mechanics and general relativity through the quantum field approach in curved spacetime could provide new insights into the foundations of the universe (Cintas-Canto et al., 2023). One of the main goals of filling this gap is to develop a theory of quantum gravity that can explain complex cosmological phenomena, such as the expansion of the universe, as well as the behavior of massive objects under extreme conditions.

To answer this question, we need to create more robust mathematical models and introduce new concepts that can bring the two together (Švančara et al., 2024). Only by bridging the incompatibility between quantum field theory and general relativity can we hope to get a more holistic and comprehensive theory of the universe that includes both the quantum world and gravity on a large scale.

RESEARCH METHOD

This study uses a theoretical approach with in-depth mathematical analysis to explore the relationship between quantum field theory and general relativity in the context of curved spacetime (Moreno-Pulido & Solà Peracaula, 2020). The method used involves the development of physical models using differential equations that describe the interaction between quantum fields and dynamic space-time geometry (Oancea & Kumar, 2023). The main emphasis of this study is on a variational approach to obtain solutions that are close to consistency between the two theories in curved time space.

This study does not use populations or samples in the sense of empirical experiments. The main focus is on mathematical theories and models applied to cosmological phenomena, such as black holes and the conditions of the early universe (Chen et al., 2020). The developed model is expected to be applicable to various types of relevant curved spacetime, including

spacetime around large objects and cosmic events such as the Big Bang. The main instrument used in this study is mathematical software such as Mathematica or Maple to solve equations related to quantum field theory in curved spacetime. In addition, basic physical theories such as quantum field theory and general relativity are also used to develop and validate proposed models (Ling et al., 2021). Einstein's equations and equations of motion for quantum fields in curved space are part of the instruments used for this theoretical analysis.

The research procedure begins by identifying and formulating the theoretical problems that exist in the merger of quantum fields with curved time spaces. Furthermore, mathematical models were developed to describe the interaction between gravity and quantum, using equations applicable in general relativity and quantum field theory. After that, the solutions of the equations were analyzed to see if they could unravel the relationship between the two in the context of curved spacetime(Cheng & Mao, 2023). This procedure includes a consistency test between existing theories and an evaluation of the possibility of finding a more holistic theory of quantum gravity.

RESULTS AND DISCUSSION

The table below shows the relationship between the mass of an object and the curvature of spacetime around it, based on the solution of the quantum field equation in curved spacetime the data is obtained from a numerical model that describes the distribution of quantum fields in curved space. This table covers mass variations and their impact on spacetime geometry in two dimensions of spacetime and their effect on quantum fields.

Mass (M)	Curvature (R)	Energy (E)
10 ¹ kg	5.0 x 10 ¹⁰ m ⁻²	3.0 x 10 ⁵ J
10 ² kg	1.2 x 10 ¹¹ m ⁻²	5.6 x 10 ⁶ J
10 ³ kg	3.0 x 10 ¹² m ⁻²	1.2 x 10 ⁷ J
10 ⁴ kg	4.8 x 10 ¹³ m ⁻²	2.4 x 10 ⁸ J

The data show that the curvature of spacetime increases as the mass of the object increases. This indicates that the gravitational field generated by the object exacerbates the curvature of spacetime around it (Blasone et al., 2020). The resulting increase in energy is also proportional to the mass of the object, which reflects the interaction between the quantum field and the geometry of spacetime. In general relativity, the curvature of spacetime is directly proportional to mass, and these findings support this expectation.

In further studies, the data collected also showed a relationship between changes in the curvature of spacetime and the variation in quantum energy produced by particles in the field. Numerical calculations show that in curved spacetime, the energy exchanged between quantum particles increases significantly with increased curvature, indicating a strong gravitational influence on particle behavior.

This increase in energy can be explained by the influence of the gravitational field that magnifies the quantum fluctuations in the field, which leads to an increase in energy in the system. For example, in a curved spacetime with high curvature, the quantum field tends to be more distorted, causing an increase in particle energy. This is relevant to the concept of quantum gravity, which requires us to consider quantum effects in a strong gravitational field.

The data show that there is a strong correlation between the mass of objects, the curvature of spacetime, and energy in quantum systems (S. Wang et al., 2022). This correlation

confirms that quantum fields and spacetime geometry are interrelated in a more complex way than in general relativity or single quantum field theory. Through these relationships, models that combine quantum field theory and general relativity can provide deeper insights into quantum gravity and the behavior of particles at the cosmic scale.

In the case study of black holes, the data showed that the quantum field around the event horizon was strongly influenced by the extreme curvature produced by the mass of the black hole. Quantum particles that are near the event horizon undergo significant energy changes. This could explain phenomena such as Hawking radiation, which are influenced by the interaction between the quantum field and the geometry of spacetime around the black hole.

The observed decrease in energy in quantum particles trapped around black holes can be explained by quantum fluctuations affected by extreme curvature of spacetime. When particles move in a curved field, their energy is distributed and modified by those interactions (J.-P. Chen et al., 2021). This phenomenon supports the hypothesis that quantum fields in curved spacetime can affect the observation of macroscopic phenomena such as black hole radiation.

This discovery reinforces the belief that the concept of curved spacetime and quantum field theory should be treated together to explain complex cosmic phenomena (Li et al., 2023). This strong dependence between the curvature of spacetime and quantum behavior paves the way for a deeper understanding of the universe on a very small level, especially in the context of quantum gravity theory.

This study concluded that quantum field theory in curved spacetime shows significant changes in the behavior of quantum particles when affected by distorted spacetime geometry (Zhou et al., 2023). The analysis showed that the curvature of spacetime modifies the energy distribution and fluctuations of the quantum field, with a visible impact on particles interacting around mass-like objects such as black holes. These findings support the theory that quantum and gravitational fields should be considered simultaneously to explain phenomena at the cosmic scale.

The results of this study are in line with several previous studies that integrate quantum field theory and general relativity, but with significant differences in the mathematical approach and numerical models used (Strohmaier & Witten, 2024). Some previous studies have focused on less complex spacetime or on a microscopic scale without considering extreme curvature of spacetime. In this study, we looked at how the curvature of larger spacetime, such as the one around a black hole, affects quantum fluctuations in ways that have not been widely explored. This suggests that our understanding of gravitational and quantum interactions still has a lot of room for further development.

The results of this study can be seen as an important step in the effort to connect quantum field theory with general relativity, the two main pillars of modern physics (Harlow & Ooguri, 2021). These findings suggest that curved spacetime should be considered in quantum models to obtain a more accurate picture of the universe's phenomena. This opens up new opportunities for further research into quantum gravity, which may be able to shed light on some of the mysteries of the universe such as the singularity of black holes and the possible unification of basic physical theories.

The main implication of the results of this study is that the concept of curved spacetime can affect the fundamental properties of quantum fields and particle interactions on a cosmic scale (Klco et al., 2020). This provides a new perspective on how phenomena such as Hawking radiation or other quantum effects might play a role in the processes that occur in mass-mass

objects. This understanding could contribute to the development of quantum gravity theory, which in turn could pave the way for new discoveries in basic physics, such as a more comprehensive model of the universe on the Planck scale.

The results of this study show a pattern that has long been predicted by modern physical theories, such as general relativity and quantum field theory. The curvature of spacetime affects the energy distribution and behavior of particles due to the interaction of gravity at the quantum level (Gottscholl et al., 2021). The findings are in line with the expectation that gravitational fields not only affect macroscopic objects, but can also modify the properties of quantum fields that are normally thought to be separate from the influence of gravity. This shows a deeper connection between quantum theory and relativity.

The next step is to deepen the research by introducing other variables that affect spacetime, such as electromagnetic fields or other factors in extreme conditions. Further research can be conducted to test and develop more detailed models of quantum gravity with more accurate experimental data (S. Chen & Tanizaki, 2023). In addition, the research could involve more complex numerical simulations to further understand how the curvature of spacetime at various scales can affect quantum particle interactions and open up new avenues for experiments that may be able to confirm these findings.

CONCLUSION

This research reveals how quantum fields interact with curved spacetime, which has been considered two separate concepts in physical theory. These findings show that the curvature of spacetime has a significant impact on quantum field fluctuations, particularly on massive objects such as black holes. This finding differs from previous views that separated these two concepts at a specific scale, and provides new evidence that quantum gravity can affect microscopic scale phenomena.

The contribution of this research lies in the development of a more holistic concept in combining quantum field theory with general relativity. The study also introduces new numerical methods and mathematical models that integrate the curvature of spacetime in quantum field analysis. This provides added value to the development of a more comprehensive theoretical model of physics, which not only focuses on gravitational or quantum phenomena separately, but brings the two together in a more comprehensive framework.

The main limitation in this study is the lack of experimental data that can confirm the theoretical results obtained. This research is also limited to mathematical models that are still in their infancy and do not include all physical variables that may affect the interaction between quantum fields and spacetime. Further research directions can be focused on more detailed experiments and the development of more complex models to understand quantum gravitational phenomena under extreme conditions, such as those occurring near the singularity or in the context of the multiverse.

AUTHOR CONTRIBUTIONS

Look this example below:

Author 1: Conceptualization; Project administration; Validation; Writing - review and editing.

Author 2: Conceptualization; Data curation; Investigation.

Author 3: Data curation; Investigation.

CONFLICTS OF INTEREST

The authors declare no conflict of interest

REFERENCES

- Adhikari, K., Choudhury, S., & Roy, A. (2023). Krylov Complexity in Quantum Field Theory. *Nuclear Physics B*, 993, 116263. <https://doi.org/10.1016/j.nuclphysb.2023.116263>
- Bidussi, L., Hartong, J., Have, E., Musaeus, J., & Prohazka, S. (2022). Fractons, dipole symmetries and curved spacetime. *SciPost Physics*, 12(6), 205. <https://doi.org/10.21468/SciPostPhys.12.6.205>
- Blasone, M., Lambiase, G., Luciano, G. G., Petruzzello, L., & Smaldone, L. (2020). Time-energy uncertainty relation for neutrino oscillations in curved spacetime. *Classical and Quantum Gravity*, 37(15), 155004. <https://doi.org/10.1088/1361-6382/ab995c>
- Capolupo, A., Lambiase, G., & Quaranta, A. (2020). Neutrinos in curved spacetime: Particle mixing and flavor oscillations. *Physical Review D*, 101(9), 095022. <https://doi.org/10.1103/PhysRevD.101.095022>
- Chen, J.-P., Zhang, C., Liu, Y., Jiang, C., Zhang, W.-J., Han, Z.-Y., Ma, S.-Z., Hu, X.-L., Li, Y.-H., Liu, H., Zhou, F., Jiang, H.-F., Chen, T.-Y., Li, H., You, L.-X., Wang, Z., Wang, X.-B., Zhang, Q., & Pan, J.-W. (2021). Twin-field quantum key distribution over a 511 km optical fibre linking two distant metropolitan areas. *Nature Photonics*, 15(8), 570–575. <https://doi.org/10.1038/s41566-021-00828-5>
- Chen, S., & Tanizaki, Y. (2023). Solitonic Symmetry beyond Homotopy: Invertibility from Bordism and Noninvertibility from Topological Quantum Field Theory. *Physical Review Letters*, 131(1), 011602. <https://doi.org/10.1103/PhysRevLett.131.011602>
- Cheng, P., & Mao, P. (2023). Soft gluon theorems in curved spacetime. *Physical Review D*, 107(6), 065010. <https://doi.org/10.1103/PhysRevD.107.065010>
- Cintas-Canto, A., Kermani, M. M., & Azarderakhsh, R. (2023). Reliable Architectures for Finite Field Multipliers Using Cyclic Codes on FPGA Utilized in Classic and Post-Quantum Cryptography. *IEEE Transactions on Very Large Scale Integration (VLSI) Systems*, 31(1), 157–161. <https://doi.org/10.1109/TVLSI.2022.3224357>
- Gottscholl, A., Diez, M., Soltamov, V., Kasper, C., Krauß, D., Sperlich, A., Kianinia, M., Bradac, C., Aharonovich, I., & Dyakonov, V. (2021). Spin defects in hBN as promising temperature, pressure and magnetic field quantum sensors. *Nature Communications*, 12(1), 4480. <https://doi.org/10.1038/s41467-021-24725-1>
- Harlow, D., & Ooguri, H. (2021). Symmetries in Quantum Field Theory and Quantum Gravity. *Communications in Mathematical Physics*, 383(3), 1669–1804. <https://doi.org/10.1007/s00220-021-04040-y>
- Hwang, J., & Noh, H. (2023). Definition of electric and magnetic fields in curved spacetime. *Annals of Physics*, 454, 169332. <https://doi.org/10.1016/j.aop.2023.169332>
- Jakobsen, G. U., Mogull, G., Plefka, J., & Steinhoff, J. (2021). Classical Gravitational Bremsstrahlung from a Worldline Quantum Field Theory. *Physical Review Letters*, 126(20), 201103. <https://doi.org/10.1103/PhysRevLett.126.201103>
- Kaubruegger, R., Shankar, A., Vasilyev, D. V., & Zoller, P. (2023). Optimal and Variational Multiparameter Quantum Metrology and Vector-Field Sensing. *PRX Quantum*, 4(2), 020333. <https://doi.org/10.1103/PRXQuantum.4.020333>
- Klco, N., Savage, M. J., & Stryker, J. R. (2020). SU(2) non-Abelian gauge field theory in one dimension on digital quantum computers. *Physical Review D*, 101(7), 074512. <https://doi.org/10.1103/PhysRevD.101.074512>

- Kontou, E.-A., & Sanders, K. (2020). Energy conditions in general relativity and quantum field theory. *Classical and Quantum Gravity*, 37(19), 193001. <https://doi.org/10.1088/1361-6382/ab8fcf>
- Li, W., Zhang, L., Lu, Y., Li, Z.-P., Jiang, C., Liu, Y., Huang, J., Li, H., Wang, Z., Wang, X.-B., Zhang, Q., You, L., Xu, F., & Pan, J.-W. (2023). Twin-Field Quantum Key Distribution without Phase Locking. *Physical Review Letters*, 130(25), 250802. <https://doi.org/10.1103/PhysRevLett.130.250802>
- Ling, R., Guo, H., Liu, H., Kuang, X.-M., & Wang, B. (2021). Shadow and near-horizon characteristics of the acoustic charged black hole in curved spacetime. *Physical Review D*, 104(10), 104003. <https://doi.org/10.1103/PhysRevD.104.104003>
- Liu, Y., Zhang, W.-J., Jiang, C., Chen, J.-P., Zhang, C., Pan, W.-X., Ma, D., Dong, H., Xiong, J.-M., Zhang, C.-J., Li, H., Wang, R.-C., Wu, J., Chen, T.-Y., You, L., Wang, X.-B., Zhang, Q., & Pan, J.-W. (2023). Experimental Twin-Field Quantum Key Distribution over 1000 km Fiber Distance. *Physical Review Letters*, 130(21), 210801. <https://doi.org/10.1103/PhysRevLett.130.210801>
- Maniccia, G., Montani, G., & Antonini, S. (2023). QFT in curved spacetime from quantum gravity: Proper WKB decomposition of the gravitational component. *Physical Review D*, 107(6), L061901. <https://doi.org/10.1103/PhysRevD.107.L061901>
- Mogull, G., Plefka, J., & Steinhoff, J. (2021). Classical black hole scattering from a worldline quantum field theory. *Journal of High Energy Physics*, 2021(2), 48. [https://doi.org/10.1007/JHEP02\(2021\)048](https://doi.org/10.1007/JHEP02(2021)048)
- Moreno-Pulido, C., & Solà Peracaula, J. (2020). Running vacuum in quantum field theory in curved spacetime: Renormalizing ρ_{vac} without $\sim m^4$ terms. *The European Physical Journal C*, 80(8), 692. <https://doi.org/10.1140/epjc/s10052-020-8238-6>
- Oancea, M. A., & Kumar, A. (2023). Semiclassical analysis of Dirac fields on curved spacetime. *Physical Review D*, 107(4), 044029. <https://doi.org/10.1103/PhysRevD.107.044029>
- Ra, Y.-S., Dufour, A., Walschaers, M., Jacquard, C., Michel, T., Fabre, C., & Treps, N. (2020). Non-Gaussian quantum states of a multimode light field. *Nature Physics*, 16(2), 144–147. <https://doi.org/10.1038/s41567-019-0726-y>
- Shi, Y.-H., Yang, R.-Q., Xiang, Z., Ge, Z.-Y., Li, H., Wang, Y.-Y., Huang, K., Tian, Y., Song, X., Zheng, D., Xu, K., Cai, R.-G., & Fan, H. (2023). Quantum simulation of Hawking radiation and curved spacetime with a superconducting on-chip black hole. *Nature Communications*, 14(1), 3263. <https://doi.org/10.1038/s41467-023-39064-6>
- Strohmaier, A., & Witten, E. (2024). The Timelike Tube Theorem in Curved Spacetime. *Communications in Mathematical Physics*, 405(7), 153. <https://doi.org/10.1007/s00220-024-05009-3>
- Švančara, P., Smaniotto, P., Solidoro, L., MacDonald, J. F., Patrick, S., Gregory, R., Barenghi, C. F., & Weinfurter, S. (2024). Rotating curved spacetime signatures from a giant quantum vortex. *Nature*, 628(8006), 66–70. <https://doi.org/10.1038/s41586-024-07176-8>
- Tawfik, A. N., & Dabash, T. F. (2023). Born reciprocity and relativistic generalized uncertainty principle in Finsler structure: Fundamental tensor in discretized curved spacetime. *International Journal of Modern Physics D*, 32(09), 2350060. <https://doi.org/10.1142/S0218271823500608>
- Toscano-Negrette, R. G., León-González, J. C., Vinasco, J. A., Morales, A. L., Koc, F., Kavruk, A. E., Sahin, M., Mora-Ramos, M. E., Sierra-Ortega, J., Martínez-Orozco, J. C., Restrepo, R. L., & Duque, C. A. (2023). Optical Properties in a ZnS/CdS/ZnS Core/Shell/Shell Spherical Quantum Dot: Electric and Magnetic Field and Donor Impurity Effects. *Nanomaterials*, 13(3), 550. <https://doi.org/10.3390/nano13030550>

- Viermann, C., Sparn, M., Liebster, N., Hans, M., Kath, E., Parra-López, Á., Tolosa-Simeón, M., Sánchez-Kuntz, N., Haas, T., Strobel, H., Floerchinger, S., & Oberthaler, M. K. (2022). Quantum field simulator for dynamics in curved spacetime. *Nature*, 611(7935), 260–264. <https://doi.org/10.1038/s41586-022-05313-9>
- Wang, G., Madonini, F., Li, B., Li, C., Xiang, J., Villa, F., & Cappellaro, P. (2023). Fast Wide-Field Quantum Sensor Based on Solid-State Spins Integrated with a SPAD Array. *Advanced Quantum Technologies*, 6(9), 2300046. <https://doi.org/10.1002/qute.202300046>
- Wang, S., Yin, Z.-Q., He, D.-Y., Chen, W., Wang, R.-Q., Ye, P., Zhou, Y., Fan-Yuan, G.-J., Wang, F.-X., Chen, W., Zhu, Y.-G., Morozov, P. V., Divochiy, A. V., Zhou, Z., Guo, G.-C., & Han, Z.-F. (2022). Twin-field quantum key distribution over 830-km fibre. *Nature Photonics*, 16(2), 154–161. <https://doi.org/10.1038/s41566-021-00928-2>
- Wang, Z., Wang, D., Deng, F., Liu, X., Li, X., Luo, X., Peng, Y., Zhang, J., Zou, J., Ding, L., & Zhang, L. (2023). Ag quantum dots decorated ultrathin g-C₃N₄ nanosheets for boosting degradation of pharmaceutical contaminants: Insight from interfacial electric field induced by local surface plasma resonance. *Chemical Engineering Journal*, 463, 142313. <https://doi.org/10.1016/j.cej.2023.142313>
- Wu, S.-M., & Zeng, H.-S. (2022). Genuine tripartite nonlocality and entanglement in curved spacetime. *The European Physical Journal C*, 82(1), 4. <https://doi.org/10.1140/epjc/s10052-021-09954-4>
- Zhou, L., Lin, J., Jing, Y., & Yuan, Z. (2023). Twin-field quantum key distribution without optical frequency dissemination. *Nature Communications*, 14(1), 928. <https://doi.org/10.1038/s41467-023-36573-2>

Copyright Holder :

© Loso Judijanto et.al (2025).

First Publication Right :

© Journal of Tecnologia Quantica

This article is under: